
TOMATO LUTADORA

LUTADORA is an exciting new determinate saladette bred in harsh environmental conditions for the Mozambique market with all the strong features needed for the tough conditions.

- Adaptable variety
- Compact, balanced plant
- Big fruit size
- Fol:3, Rs, TMV and TSWV resistance





A determinate saladette tomato suited to open field conditions.

Maturity

It is an early-medium maturing variety. The first fruit will mature around 75-80 days after transplanting. Days to maturity may vary due to environmental conditions that may delay maturity.

Plant Characteristics

LUTADORA is compact, with a strong stand and plant frame, making the variety balanced and well adapted to harsh conditions.

Plant Populations

Depending on spacing the variety can be planted between 13500 – 15000 plants per hectare. It is an open field variety requiring one to two wires to help keep it upright with the fruit set.

Variety Characteristics

LUTADORA has firm, large, attractive, egg-shaped fruit with an average mass of 120-140 g. It is well balanced and keeps its fruit shape and size.

Features & Benefits

LUTADORA has an excellent disease resistance package, with good fruit size in comparison to Star 9068.

Disease Resistance

HR: Va:1, Vd:1, Fol: 1-3, TMV; IR: Ma, Mi, Mj, Rs, For, Lt, TSWV

Disclaimer: This information is based on our observations and/or information from other sources. As crop performance depends on the interaction between the genetic potential of the seed, its physiological characteristics, and the environment, including management, we give no warranty express or implied, for the performance of crops relative to the information given nor do we accept any liability for any loss, direct or consequential, that may arise from whatsoever cause. Please read the Starke Ayres Standard Terms and Conditions of Sale before ordering seed.

Resistance: is the ability of a plant variety to restrict the growth and development of a specified pest or pathogen and/or the damage they cause when compared to susceptible plant varieties under similar environmental conditions and pest or pathogen pressure. Resistant varieties may exhibit some disease symptoms or damage under heavy pest or pathogen pressure (HR = High resistance, IR = Intermediate resistance).